arXiv:1108.0601v2 [math.AT] 13 Nov 2011

A FREE FROBENIUS BIALGEBRA ON THE DIFFERENTIAL FORMS OF A MANIFOLD

MICAH MILLER

Abstract. Let M be a closed, oriented manifold. We prove that the quasi-isomorphism class of the $Frob^0_\infty$-bialgebra structure on $H^*(M)$ induced by the open TFT on $\Omega^*(M)$ is a homotopy invariant of the manifold. This is a three step process. First, we describe the $Frob^0_\infty$-bialgebra on $H^*(M)$ induced by the partial $Frob^0_\infty$-bialgebra on $\Omega^*(M)$. We then describe the $Frob^0_\infty$-bialgebra on $H^*(M)$ induced by the cyclic C_∞-algebra on $H^*(M)$. Finally, we show these two $Frob^0_\infty$-bialgebras are the same. Since the cyclic C_∞-algebra is a homotopy invariant, this proves our claim.

Contents

1. Introduction 1
2. Operads and dioperads 3
2.1. Cyclic operads 3
2.2. Dioperads 4
2.3. Frobenius dioperad 7
2.4. Functor from cyclic operads to dioperads 7
2.5. Algebras over cyclic operads and dioperads 9
2.6. Partial algebras over cyclic operads and dioperads 10
3. Transfer of structure 10
3.1. Transferring A_∞ and C_∞-algebras by a contraction 10
3.2. Transfer of partial $Frob^0_\infty$-bialgebra by a contraction 11
4. Topological conformal field theories 13
4.1. Cellular differential forms 15
4.2. Subspaces of the moduli space 17
5. Constructing a partial $Frob^0_\infty$-bialgebra on $Curr_*(M)$ 17
5.1. Double forms $\Omega^*(M \times M)$ 17
5.2. Domain for $Curr_*(M)$ 18
5.3. Partial $Frob^0_\infty$-bialgebra structure on $Curr_*(M)$ 19
5.4. Sub partial $Frob^0_\infty$-bialgebra on $\Omega^*(M)$ 21
6. Homotopy invariance of the $Frob^0_\infty$-bialgebra 22
References 23

1. Introduction

In this paper, we investigate the algebraic structure on the differential forms of a closed, oriented manifold $\Omega^*(M)$ and the induced structure on its cohomology

$H^*(M)$. It is well known that $H^*(M)$ is a Frobenius abilgebra, i.e., there is a multiplication and comultiplication, such that the comultiplication is a map of modules. The structure on $\Omega^*(M)$ which induce Frobenius bialgebra is less well understood. We provide a model for this structure and show that when we transfer the maps to $H^*(M)$, we obtain a $Frob^0_\infty$-bialgebra. We then show that this $Frob^0_\infty$-bialgebra is a homotopy invariant of the manifold.

To do this, we recall in Section 2 the definition and properties of operads and dioperads. The main examples in this paper are

(1) the cyclic C_∞ operad, which is used to describe the cyclic C_∞-algebra on $H^*(M)$ induced by the wedge product and Poincare Duality,
(2) the $Frob^0_\infty$ dioperad, which is used to describe the Frobenius bialgebra on $H^*(M)$,

We will also need to discuss the notion of partial algebras. These objects were discussed by Wilson, [W], in the context of operads. We extend the discussion to include dioperads.

In Section 3, we discuss methods of transferring algebraic structures from a chain complex to its homology as a way to build minimal models. Kontsevich and Soibelman show, given a contraction between an A_∞-algebra $(A_*, \{m_n^A\})$ and a chain complex B_*, how to define an A_∞-algebra structure $\{m_n^B\}$ on B_*, [KS]. Cheng and Getzler, [CG], show that if A_* is a C_∞-algebra, then $\{m_n^B\}$ will be a C_∞-algebra structure. In ([**?**], Appendix B), the authors provide an explicit way to transfer a P_∞-algebra structure for an operad P. Following these methods, we describe how to transfer a $Frob^0_\infty$-algebra.

With the abstract setting in place, the rest of the paper focuses on constructing a partial $Frob^0_\infty$-bialgebra structure on $\Omega^*(M)$ in two ways. First, we consider the partial Frobenius bialgebra on $Curr_*(M)$, using ideas from topological conformal field theory. This construction makes use of the heat kernel $\widehat{K}_t : Curr_*(M) \to Curr_*(M)$, which can be thought of as a contraction between $Curr_*(M)$ and itself. The differential forms $\Omega^*(M)$ sit inside $Curr_*(M)$ as smooth currents, and we show that the partial $Frob^0_\infty$-bialgebra restricts to $\Omega^*(M)$. We apply the transfer theorem to obtain a $Frob^0_\infty$-bialgebra structure on$H^*(M)$.

The second approach for constructing a $Frob^0_\infty$-bialgebra is to define a functor

$$\text{cyclic operads} \to \text{dioperads}$$

to the map of cyclic operads C_∞operad $\to End(H^*(M))$ to obtain $Frob^0_\infty \to End(H^*(M))$. This will also be a $Frob^0_\infty$-bialgebra. We show in Section 6 that this $Frob^0_\infty$-bialgebra is the same as the previous one. Since $(H^*(M), \{m_n\}, PD)$ is known to be a homotopy invariant, this proves the claim.

Remark 1.1. A note on terminology is in order. A Frobenius algebra consists of a vector space V with a commutative multiplication and a non-degenerate bilinear form $\langle , \rangle$, such that $\langle a, bc\rangle = \langle ab, c\rangle$. Using the language of operads, a Frobenius algebra is an algebra over the cyclic commutative operad.

A Frobenius bialgebra consists of a vector space V with a commutative multiplication and cocommutative comultiplication $\Delta : V \to V \otimes V$, such that Δ is a map of bimodules. That is,

$$\Delta(ab) = a_{(1)} \otimes (a_{(2)} b) = (ab_{(1)}) \otimes b_{(2)}.$$

Frobenius bialgebras arise in the study topological quantum field theories. They are related to Frobenius algebras and sometimes referred to as Frobenius algebras. Since we discuss both structures, we will use Frobenius algebras and Frobenius bialgebras to make the distinction.

Acknowledgements: This paper would not have been possible without the help and direction of Mahmoud Zeinalian. Scott Wilson has also been generous with his comments and ideas on the paper, which have helped in improving the paper. The author is also grateful for conversations with Józef Dodziuk and Dennis Sullivan concerning topics covered in this paper.

2. Operads and dioperads

2.1. **Cyclic operads.** We first review the definition of an operad. Let $\mathcal{C}$ be the category of $\mathbb{Z}$ graded vector spaces over a field k of characteristic zero.

Definition 2.1. An **operad** P in $\mathcal{C}$ consists of

(1) objects $P(n)$ in $\mathcal{C}$ with $\mathbb{S}_n$ action for $n > 0$
(2) for $m, n > 0$ and $1 \leq i \leq m$, morphisms
$$\circ_i : P(m) \otimes P(n) \to P(m+n-1),$$
(3) a morphism $\eta : k \to P(1)$

satisfying the following associativity and equivariance conditions,

(1) For compositions of $P(m) \otimes P(n) \otimes P(p)$, the following associative condition must be satisfied:
$$\circ_i(\circ_j) \otimes 1) = \begin{cases} \circ_{j+p-1}(\circ_i \otimes 1)(1 \otimes \tau), & \text{for } 1 \leq i < j-1, \\ \circ_j(1 \otimes \circ_{i-j+1}), & \text{for } j \leq i \leq j+n-1, \\ \circ_j(\circ_{i-n+1} \otimes 1)(1 \otimes \tau), & \text{for} j+n \leq i, \end{cases}$$
where $\tau : P(n) \otimes P(p) \to P(p) \otimes P(n)$ is the transposition given by the symmetry,
(2) the operation is $\circ_i$ is equivariant in the sense that
$$\circ_i(\sigma\rho) = (\sigma \circ_i \rho)\circ_{\sigma(i)}$$
on $P(m) \otimes P(n)$ where $\sigma \in S(m)$, $\rho \in S(n)$, and $\sigma \circ_i \rho$ is a block permutation

A cyclic operad is an operad P such that the $\mathbb{S}_n$ action on $P(n)$ extends to an action on $\mathbb{S}_n^+$ satisfying associative and equivariance relations. Here, $\mathbb{S}_n^+$ is the permutation group on $\{0, 1, 2, \cdots, n\}$. The group $\mathbb{S}_n$ is viewed as the subgroup of permutations in $\mathbb{S}_n^+$ that fix 0. Let τ_n be the cyclic permutation $(0, 1, \cdots, n)$. Then an operad P is cyclic if the $\mathbb{S}_n^+$ action satisfies the following.

(1) If $\eta : 1 \to P(1)$ is the unit, then the following diagram commutes

$$\begin{tikzcd} P(1) \arrow[r, "\tau_1"] & P(1) \\ 1 \arrow[u, "\eta"] \arrow[ur, "\eta"'] & \end{tikzcd}$$

(2) For each $m, n \geq 1$, the following diagram commutes

$$\begin{tikzcd}
P(m) \otimes P(n) \arrow[r, "\circ_1"] \arrow[d, "\tau_m \otimes \tau_n"] & P(m+n-1) \arrow[dd, "\tau_{m+n-1}"] \\
P(m) \otimes P(n) \arrow[d] & \\
P(n) \otimes P(m) \arrow[r, "\circ_n"] & P(m+n-1)
\end{tikzcd}$$

If V is a finite dimensional vector space with a non-degenerate bilinear form $\langle , \rangle$, then $End(V)$ is a cyclic operad. The associative, commutative, and Lie operad are examples of cyclic operads, as shown in [GK].

2.2. **Dioperads.** Dioperads model structures on vector spaces with products and coproducts, where the composition rule glues one output to one input. In this way, genus is never created. Dioperads can model Frobenius bialgebras, since the relation that the coproduct be a bimodule map does not have genus and can be described by grafting one output of a tree with one input of another tree. In contrast, dioperads cannot model bialgebras, since the requirement that the coproduct be an algebra map introduces genus.

Definition 2.2. A **dioperad** P in $\mathcal{C}$ consists of

1. objects $P(m, n)$ in $\mathcal{C}$ with an $(\mathbb{S}_m, \mathbb{S}_n)$ bimodule structure for $m, n > 0$,
2. for $m_1, m_2, n_1, n_2 \geq 1$, $1 \leq i \leq n_1$, and $1 \leq j \leq m_2$, morphisms
$$_i\circ_j : P(m_1, n_1) \otimes P(m_2, n_2) \to P(m_1 + m_2 - 1, n_1 + n_2 - 1),$$
3. a morphism $\eta : k \to P(1, 1)$ such that
$$_1 \circ_i (\eta \otimes Id) : k \otimes P(m, n) \tilde{\to} P(m, n),$$
and
$$_j \circ_1 (Id \otimes \eta) : P(m, n)) \tilde{\to} P(m, n)$$
are the canonical isomorphisms.

The morphisms $_i\circ_j$ must respect the $(\mathbb{S}_m, \mathbb{S}_n)$ bimodule actions. The conditions are described in [G].

The augmentation ideal of a dioperad P, denoted $\overline{P}$, is defined by $\overline{P}(1, 1) = 0$ and $\overline{P}(m, n) = P(m, n)$ for $m + n \geq 3$.

If V is an object in $\mathcal{C}$, the *endomorphism dioperad* $End^0(V)$ is given by letting $End^0(V)(m, n)$ be the complex $Hom(V^{\otimes n}, V^{\otimes m})$. The superscript 0 indicates that only maps without genus are considered in the complex $Hom(V^{\otimes n}, V^{\otimes m})$ and distinguishes the endomorphism dioperad $End^0(V)$ from the endomorphism properad $End(V)$. An algebra over a dioperad P is a dioperad map $P \to End^0(V)$.

If V is an $(\mathbb{S}_m, \mathbb{S}_n)$ bimodule, V^* is also an $(\mathbb{S}_m, \mathbb{S}_n)$ bimodule. If $f \in V^*$ and $\sigma \in \mathbb{S}_m$, then $\sigma f \in V^*$ is defined by $\sigma f(x) = f(\sigma(x))$. The $\mathbb{S}_n$ action on V^* is defined similarly.

There is an equivalent definition of a dioperad as a monoid in a monoidal category. We describe the category of two leveled trees. A tree will be a finite, directed tree, with m inputs and n outputs. If every vertex has valence at least three, then T is said to be reduced. Each tree T has a global flow- a direction on the edges signifying



Figure 2.1. A 2 leveled tree.



Figure 2.2. The equivalence relation in the definition $\boxtimes$.

inputs and outputs. Let $Edge(T)$ be the set of edges of T, $edge(T)$ be the set of internal edges of T, $In(v)$ be the set of incoming edges at the vertex $v \in T$, and $Out(v)$ be the set of outgoing edges at v. We assume that $In(v)$ and $Out(v)$ are labelled by $\{1, 2, \cdots, |In(v)|\}$ and $\{1, 2, \cdots, |Out(v)|\}$. For a k- dimensional vector space V, denote by $Det(V)$ be the top exterior power of V. The global inputs and outputs will also be labelled. A 2-*leveled tree* is a directed tree such that the vertices are divided into a top level and bottom level. Denote the set of vertices in the top level by N_1 and bottom level by N_2. We refer the reader to Figure 2.1. We let $\mathcal{T}$ be the category of graphs and $\mathcal{T}^2$ be the category of 2 leveled trees.

If $\{P(m, n)\}$ and $\{Q(m, n)\}$ are collections of two $(\mathbb{S}_m, \mathbb{S}_n)$-bimodules, their composition product is given by considering the collection of two leveled trees, identifying vertices of the top level with elements in $P(m, n)$ and vertices in the bottom level with elements in $Q(m, n)$. The composition product $\boxtimes$ of P and Q is given by

$$Q \boxtimes P = \left(\oplus_{g \in \mathcal{T}^2} \otimes_{v \in N_2} Q(|Out(v), In(v)) \otimes_k \otimes_{v \in N_1} P(|Out(v)|, |In(v)|) \right) / \sim,$$

where the equivalence is explained in Figure 2.2 .

The connected composition product $\boxtimes_c$ is obtained by restricting $\boxtimes$ to connected two leveled trees. The unit I for this monoidal product is

$$I(m, n) := \begin{cases} \mathbb{K} & \text{for } (m, n) = (1, 1) \\ 0 & \text{otherwise.} \end{cases}$$

In [G], another product is defined on the category of $\mathbb{S}$-bimodules, denoted $\boxdot$. It is obtained from $\boxtimes_c$ by restricting over saturated two level trees. A vertex $v \in T$ is saturated if every path from a leaf vertex to a root vertex passes through v. A tree is called saturated if every vertex is saturated. Then if P_1 and P_2 are two $\mathbb{S}$ bimodules, define $P_1 \boxdot P_2$ by

$$(P_1 \boxdot P_2)(m, n) = \bigoplus_{(T,l)} \otimes_{i=1}^{2} \otimes_{v \in l^{-1}} P_i(Out(v), In(v)), \tag{2.1}$$

where the sum is taken over all saturated 2 level labeled (m, n) trees (T, l). This product has a nice property with respect to Koszul duality, which we describe in Proposition 2.4 after defining the cobar complex.

Example 2.3. We describe the free dioperad. Let $E(m, n)$ is a collection of $(\mathbb{S}_m, \mathbb{S}_n)$ bimodules. Then for a tree T, let $E(T) = \otimes_{v \in T} E(Out(v), In(v))$. The free dioperad $\mathcal{F}(E)$ generated by E is given by

$$\mathcal{F}(E)(m,n) = \bigoplus_{(m,n) \text{ trees}} E(T)/\sim,$$

where the equivalence relation is the one described in the composition product $\boxtimes$ (see Figure 2.2).

Suppose $\{E(m, n)\}$ is an $(\mathbb{S}_m, \mathbb{S}_n)$ bimodule, such that $E(m, n) = 0$ for all (m, n), except $E(2, 1)$ and $E(1, 2)$. Take the free dioperad generated by E. Let (R) be an ideal generated by $R(1, 3) \subset \mathcal{F}(E)(1, 3)$, $R(3, 1) \subset \mathcal{F}(E)(3, 1)$, and $R(2, 2) \subset \mathcal{F}(2, 2)$. Then the quotient dioperad $\mathcal{F}(E)/(R)$, denoted $\langle E; R\rangle$, is called a quadratic dioperad.

Let $P = \langle E; R\rangle$ be a quadratic operad. The quadratic dual of P is a dioperad $P^! = \langle E^\vee; R^\perp\rangle$, where $E^\vee(m, n) = E^* \otimes (Sgn_m \otimes Sgn_n)$ and $R^\perp(m, n)$ is the orthogonal complement of $R(m, n)$.

Let $Det(T)$ be the one dimensional vector space $Det(k^{Edge(T)})$ and $det(T) = Det(k^{edge(T)})$. Given a dioperad P, the *cobar complex* of $P(m, n)$, denoted $C(P)$ is a bicomplex $\mathcal{F}(\overline{P}^*(m, n))$ where the differential is the sum of the differential induced by the differential of P, denoted d', and the differential induced by edge contractions, denoted d''. Looking at d'' differential, we get a complex

$$\overline{P}(m,n)^* \xrightarrow{d''} \bigoplus_{\substack{|edge(T)|=1 \\ \text{reduced } (m,n) \text{ trees}}} \overline{P}^*(T) \otimes det(T) \xrightarrow{d''} \bigoplus_{\substack{|edge(T)|=2 \\ \text{reduced } (m,n) \text{ trees}}} \overline{P}^*(T) \otimes det(T) \xrightarrow{d''} \cdots,$$

where $\overline{P}(m, n)^*$ is placed in degree one.

The *cobar dual* $\mathbb{D}(P)$ is given by $\Lambda^{-1}(C(P))$. The chain complex with d'' is given by

$$\overline{P}(m,n)^\vee \xrightarrow{d''} \bigoplus_{\substack{|edge(T)|=1 \\ \text{reduced } (m,n) \text{ trees}}} \overline{P}^*(T) \otimes Det(T) \xrightarrow{d''} \bigoplus_{\substack{|edge(T)|=2 \\ \text{reduced } (m,n) \text{ trees}}} \overline{P}^*(T) \otimes Det(T) \xrightarrow{d''} \cdots,$$

where $\overline{P}(m, n)^\vee$ is put in degree $3 - m - n$. The bimodule $\overline{P}(m, n)^\vee$ is $\overline{P}^* \otimes (Sgn_m \otimes Sgn_n)$.

Recall that a dioperad P is Koszul if there exists a dioperad Q such that $\mathbb{D}(Q) \to P$ is a quasimorphism. The dioperad Q is said to be Koszul dual to P, and is denoted by $P^!$.

Proposition 2.4 ([G], Proposition 5.9)**.** *Let $P = \langle E; R\rangle$ be a quadratic dioperad $A(n) := P(n, 1)$ and $B(n) := P^{op}(n, 1)$.*

1. *If P is a Koszul dioperad and $P(m, n) = A \boxdot B^{op}(m, n)$ for $(m, n) = (2, 2)$, $(2, 3)$, and $(3, 2)$, then $P = A \boxdot B^{op}$. Similarly, if $P(m, n) = A^{op} \boxdot B(m, n)$ for $(m, n) = (2, 2)$, $(2, 3)$ and $(3, 2)$, then $P = A^{op} \boxdot B$.*
2. *$P = A \boxdot B^{op}$ if and only if $P^! = B^{!op} \boxdot A^!$. Similarly, $P = A^{op} \boxdot B$ if and only if $P^! = B^! \boxdot A^{!op}$*

(3) *If A and B are Koszul, and $P = A \boxdot B^{op}$, then P is Koszul. Similarly, if A and B are Koszul and $P = A^{op} \boxdot B$, then P is Koszul.*

2.3. **Frobenius dioperad.** The Frobenius dioperad, denoted $Frob^0$ is generated by an element in $m \in E(2,1)$ and $\Delta \in E(1,2)$, each with trivial $\mathbb{S}_2$ actions. The generators are required to satisfy the following relations:

$$\begin{gathered}(m {}_1\circ_1 m) - (m {}_1\circ_1 m)\sigma, (m {}_1\circ_1 m) - (m {}_1\circ_1 m)\sigma^2 \\ (\Delta {}_1\circ_1 \Delta) - \sigma(\Delta {}_1\circ_1 \Delta), (\Delta {}_1\circ_1 \Delta) - \sigma^2(\Delta {}_1\circ_1 \Delta) \\ (\Delta {}_1\circ_1 m) - (m {}_1\circ_1 \Delta), (\Delta {}_1\circ_1 m) - (m {}_1\circ_2 \Delta), \\ (\Delta {}_1\circ_1 m) - (m {}_2\circ_1 \Delta), (\Delta {}_1\circ_1 m) - (m {}_2\circ_2 \Delta),\end{gathered}$$

where $\sigma = (123) \in \mathbb{S}_3$. The superscript in $Frob^0$, which we use to distinguish it from properads, emphasizes that we are considering only genus zero operations. Note that $Frob^0(m,n)$ is one dimensional for each m and n.

From Proposition 2.4, we can see the following equality: $Frob^0 = Com^{op} \boxdot Com$. It is then Koszul, with Koszul dual $Lie \boxdot Lie^{op}$. Algebras over $Lie \boxdot Lie^{op}$ are Lie bialgebras, so we denote the dioperad by $BiLie^0$. The dioperad $BiLie^0$ is generated by elements $l \in E(2,1)$ and $\delta \in E(1,2)$ using the sign action of $\mathbb{S}_2$. Its relations are spanned by

$$\begin{gathered}(l {}_1\circ_1 l) + (l {}_1\circ_1 l)\sigma + (l {}_1\circ_1)\sigma^2 \\ (\delta {}_1\circ_1 \delta) + \sigma(\delta {}_1\circ_1 \delta) + \sigma^2(\delta {}_1\circ_1 \delta) \\ \delta {}_1\circ_1 l) - (l {}_1\circ_1 \delta) - (l {}_1\circ_2 \delta) - (l {}_2\circ_1 \delta) - (l {}_2\circ_2 \delta).\end{gathered}$$

2.4. **Functor from cyclic operads to dioperads.** Given a cyclic operad P, we show how to obtain a dioperad denoted $\mathcal{D}(P)$. To see why such a construction is possible, we consider the case of the cyclic endomorphism operad $End_{cyc}(V)$. If V is a finite dimensional vector space with a non-degenerate bilinear form $\langle , \rangle : V \otimes V \to k$, then $End(V)$ is a cylic operad. The $\mathbb{S}_m^+$ action is given by viewing a map $f : V^{\otimes m} \to V$ as

$$\begin{array}{rcl}\hat{f} : V^{\otimes m+1} & \to & k \\ (v_0, \cdots, v_m) & \mapsto & \langle v_0, f(v_1, \cdots, v_m)\rangle.\end{array}$$

If f is viewed graphically as a tree with m whose leaves are labeled $1, \cdots, m$ and a root, then $\hat{f}$ is the same tree, with the root labeled by 0.

The maps $\circ_j : End(V)(m) \otimes End(V)(n) \to End(V)(m+n-1)$ are defined by grafting the root of one tree to the edge with label j to the other tree. For $\hat{f} : V^m \to k$ and $\hat{g} : V^n \to k$, if $j \neq 0$, let $\hat{f} \circ_i \hat{g}$ be $\widehat{f \circ_i g}$; that is,

$$\begin{array}{rcl}\hat{f} \circ_i \hat{g} : V^{m+n} & \to & k \\ (v_0, \cdots, v_{m+n-1}) & \to & \langle v_0, f \circ_i g(v_1, \cdots, v_{m+n-1})\rangle.\end{array}$$

If the grafting occurs along a root edge, we define the composition rule as follows,

$$\begin{array}{rcl}\hat{f} \circ_0 \hat{g} : V^{m+n} & \to & k \\ (v_0, \cdots, v_{m+n-1}) & \mapsto & \langle v_0, g(f(v_1, \cdots, v_m), v_{m+1}, \cdots, v_{m+n-1})\rangle.\end{array}$$

The motivation for the definition can be seen from drawing the tree obtained by grafting to the root edge; we refer the reader to Figure 2.3.

To this cyclic operad, we associated a dioperad denoted $\mathcal{D}(End(V))$. Let $\mathcal{D}(End(V))(m,n) = End_{cyc}(V)(m+n-1)$. The $(\mathbb{S}_m, \mathbb{S}_n)$ bimodule structure of $\mathcal{D}(End(V))(m,n)$ is given



Figure 2.3. The result of grafting two trees along their root edges. Note that the final leaf, $m+n$ is relablled zero and designated to be the root.

by viewing $\mathbb{S}_m$ and $\mathbb{S}_n$ as subgroups of $\mathbb{S}^+_{m+n-1}$. The group $\mathbb{S}_m$ is viewed as a subgroup of $\mathbb{S}^+_{m+n-1}$ by the usual identification (i.e. $1 \mapsto 1, 2 \mapsto 2, \cdots, m \mapsto m$). The group $\mathbb{S}_n$ is a subgroup of $\mathbb{S}^+_{m+n-1}$ given by the identification $1 \mapsto m+1, 2 \mapsto m+2, \cdots, n-1 \mapsto m+n-1, n \mapsto 0$.

To define the ${}_i\circ_j$ maps, we use the maps $\hat{f} \circ_j \hat{g}$ along with the τ_n. If f is a tree with m_1 inputs and n_1 outputs g is a tree with m_2 inputs and n_2 outputs, then $f\,{}_i\circ_j\, g$ is a tree with $m_1 + m_2 - 1$ inputs and $n_1 + n_2 - 1$ outputs obtained by grafting the i^{th} output of g with the j^{th} input of f. For $\hat{f}$ and $\hat{g}$, where the trees have only one output, we interpret the i^{th} output of $\hat{g}$ to be the root of $\hat{g}$ after τ^i_n has been applied to $\hat{g}$. We then apply τ^{-i}_n to undo the application. In summary, $\hat{f}\,{}_i\circ_j\,\hat{g} = \tau^{-i}_n(\hat{f} \circ_j \tau^i_n(\hat{g}))$.

This construction works for general cyclic operad, defining a functor

$$\mathcal{D} : \{\text{cyclic operads}\} \to \{\text{dioperads}\}.$$

Let P be a cyclic operad. Then define $\mathcal{D}(P)(m,n)$ to be the object $P(m+n-1)$. The $(\mathbb{S}_m, \mathbb{S}_n)$ bimodule structure is inherited from the $\mathbb{S}^+_{n+m-1}$ structure on $P(m+n-1)$, using the same identifications as above to view $\mathbb{S}_m$ and $\mathbb{S}_n$ as subgroups of $\mathbb{S}^+_{m+n-1}$. Since τ^i and $\circ_j$ maps are defined for any cyclic operad, the ${}_i\circ_j$ maps can be defined in the same way as above.

If we have a cyclic operad morphism $f : P \to Q$, there is a corresponding dioperad morphism $\mathcal{D}(P) \to \mathcal{D}(Q)$. The morphism is defined by sending $\mathcal{D}(P)(m,n) = P(m+n-1)$ to $f(P(m+n-1)) \subset Q(m+n-1) = \mathcal{D}(Q)(m,n)$.

Proposition 2.5. *There exists an exact functor*

$$\{\textit{cyclic operads}\} \to \{\textit{dioperads}\},$$

which sends the cyclic endomorphism operad to the endomorphism dioperad.

Proof. The functor is defined above. It is an exact functor because given an exact sequence

$$\cdots \to P_{i+1} \to P_i \to P_{i-1} \to \cdots,$$

the terms in the induced sequence

$$\cdots \to \mathcal{D}(P_{i+1}) \to \mathcal{D}(P_i) \to \mathcal{D}(P_{i-1}) \to \cdots$$

are the same. The difference between P and $\mathcal{D}(P)$ lies in the actions and gluing compositions, but this is not used for exactness. So the induced sequence is exact as well.

For the claim about the cyclic endomorphism operad going to the endomorphism dioperad, note that $\mathcal{D}(End(V))(m,n)$ is the bimodule $Hom(V^{\otimes m+n-1}, V) \cong V^{\otimes n+m}$. Using the isomorphism between V and V^*, we can write this as $V^{*\otimes m} \otimes$

$V^n \cong Hom(V^{\otimes m}, V^{\otimes n})$, so $\mathcal{D}(End(V))$ is at least a sub-dioperad of the endomorphism dioperad.

Again using Hom-Tensor adjunction and the isomorphism given by $\langle , \rangle$, an element in $Hom(V^{\otimes m}, V^{\otimes n})$ can be written as an element in $Hom(V^{\otimes m+n-1}, V)$. So $\mathcal{D}(End(V))(m,n)$ has all elements of the $(\mathbb{S}_m, \mathbb{S}_n)$ bimodule of the endomorphism dioperad $End(V)(m,n)$. □

Corollary 2.6. *Let V be a finite dimensional vector space. A $Frob_\infty$-algebra structure on V defines a $Frob^0_\infty$-bialgebra structure on V.*

Proof. It is known that a unital Frobenius algebra (that is, a unital cyclic commutative algebra) is the same as a unital, counital Frobenius bialgebra, (see for instance [A]). One direction can be seen by proving that the cyclic commutative operad goes to the Frobenius dioperad.

Let Com be the commutative operad, where $Com(n)$ is the one dimensional space with trivial $\mathbb{S}_n$ action. This operad is generated by $Com(2)$ with the relation that all ways of repeatedly gluing $Com(2)$ with itself are equivalent. We see that $\mathcal{D}(Com)$ is generated by $\mathcal{D}(Com(2,1)$ and $\mathcal{D}(Comm)(1,2)$, with the relation that any way of obtaining an element in $\mathcal{D}(Com)(m,n)$ through gluing generators are equivalent. The Frobenius dioperad is described in the same way, so $\mathcal{D}(Com) = Frob^0$.

A C_∞ operad, denoted $(\Gamma(Com), \partial)$ is a cofibrant model for Com; that is, $\Gamma(Com)$ is a free operad, with a differential such that $(\Gamma(Com), \partial) \to Com$ is a quasi-isomorphism. Since $\mathcal{D}$ is an exact functor, when we apply $\mathcal{D}$ to $(\Gamma(Com), \partial) \to Com$, we obtain a quasi-isomophism $\mathcal{D}(\Gamma(Com), \partial) \to Frob^0$.

To show that $\mathcal{D}(\Gamma(Com))$ is a free dioperad, we use $\mathbb{D}\mathbb{D}(Com)$ as a model for $\Gamma(Com)$, where $\mathbb{D}$ is the cobar dual functor. We claim $\mathcal{D}(\mathbb{D}(Com))$ is isomorphic to $\mathbb{D}(\mathcal{D}(Com)) = \mathbb{D}(Frob^0)$.

□

2.5. **Algebras over cyclic operads and dioperads.** As mentioned earlier, operads and dioperads are used to model algebraic structures on vector spaces. If P is a cyclic operad or dioperad, then V is a P-algebra if there is a map of cyclic operads or dioperads

$$P \to End(V).$$

Let P be a quadratic dg operad or dioperad and $P^!$ be its quadratic dual. Then P is Koszul if $\mathbb{D}(P^!) \to P$ is a quasi-isomorphism. The interest in $\mathbb{D}(P^!)$ arises from the fact that P_∞-algebras are algebras over $\mathbb{D}(P^!)$. That is, a chain complex A_* is a P_∞-algebra if there exists a map of dg operads

$$\mathbb{D}(P^!) \to End(A_*).$$

Markl, in [M], shows that algebras over resolutions have properties that one would expect a homotopy algebra to satisfy. For example, if a P_∞-algebra A is chain equivalent to a chain complex B, then B is a P_∞-algebra and the chain maps in the equivalence extend to P_∞-algebra morphisms. Sullivan, in [S], discusses generalizations of this feature to include dioperads and properads.

We define a functor functors

$$\mathcal{D} : \text{cyclic operads} \to \text{dioperads}$$

which will be used to construct $Frob^0_\infty$-algebras. This functor $\mathcal{D}$ is exact, so it sends resolutions to resolutions. Furthermore, if $\mathbb{D}(P^!) \to End^{cyc}(A_*)$ is a cyclic

P_∞-algebra, then applying $\mathcal{D}$ to the cyclic operad map results in a map of dioperads $\mathcal{D}(\mathbb{D}(P^!)) \to End^0(A_*)$. Since $\mathcal{D}(\mathbb{D}(P^!))$ is a cofibrant resolution for the dioperad $\mathcal{D}(P)$, the dioperad map defines a $\mathcal{D}(P)_\infty$-algebra structure on A_*.

2.6. **Partial algebras over cyclic operads and dioperads.** We will be interested in situations when the algebraic operations on the vector space V have domains or ranges that are not $V^{\otimes n}$. These situations arise when looking at a coproduct on forms $\Delta_t : \Omega^*(M) \to \Omega^*(M \times M)$ or product on currents $\wedge_t : Curr_*(M \times M) \to Curr_*(M)$. We will call such objects partial algebra (coalgebra) structures, although there is a difference between these objects and the partial algebras found in [W].

Definition 2.7. A **domain** for a cochain complex (A, d) is a sequence of complexes $\{A_i\}$ such that

1. $A_1 = A$,
2. $A^{\otimes i}$ is a subcomplex of A_i,
3. there is an $\mathbb{S}_i$ action on A_i, such that for $i_1 + \cdots + i_k = i$, $A^{\otimes i_1} \otimes \cdots \otimes A^{\otimes i_k}$ is an $\mathbb{S}_i$ invariant subspace of A_i,
4. $\mathbb{S}_i$ equivariant maps $A^{\otimes i} \hookrightarrow A_i$ and $A_i \to A^{\otimes i}$ whose compositions induce the identity map on cohomology,

Given a domain $\{A_i\}$ for (A, d), we can form partial endomorphism operad or dioperad. The definition of domain so far includes an $\mathbb{S}_n$ action on A_n. Let $\underline{End}(A)$ be the bimodule $\underline{End}(A)(m, n) = Hom(A_m, A_n)$. Given any $f : A_i \to A_j$, we have a map

$$\widehat{f} : A^{\otimes i} \hookrightarrow A_i \xrightarrow{f} A_j \to A^{\otimes j}.$$

So if $End(A)$ is an operad dioperad, $\underline{End}(A)$ is an operad or dioperad using the maps in (4) as above to define the composition maps. For example, for $f : A_{i_1} \to A_{j_1}$ and $g : A_{i_2} \to A_{j_2}$, let $f\ {}_i{\circ}_j\ g$ be defined by

$$A_{i_1+i_2-1} \to A^{\otimes i_1+i_2-1} \xrightarrow{\widehat{f}\ {}_i{\circ}_j \widehat{g}} A^{\otimes j_1+j_2-1} \to A_{j_1+j_2-1}.$$

We focus mainly on the partial endomorphism dioperad. The partial endomorphism dioperad is denoted $\underline{End}(A)^0(m, n) = Hom(A_m, A_n)$, (where the superscript 0 indicates only genus zero operations are considered). If P is a dg dioperad, then A is a partial P-algebra if there is a map of dg dioperads, $P \to \underline{End}^0(A)$.

Remark 2.8. This definition of domain differs from the one found in [W]. In there, each A_i is a subcomplex of $A^{\otimes i}$, and the main example is the subcomplex of transversally intersecting chains in $C_*(M)$. The difference is mostly formal and the methods used in one situation translates in an obvious way to the other situation.

3. Transfer of structure

3.1. **Transferring A_∞ and C_∞-algebras by a contraction.** Let A and B be cochain complexes over a field of characteristic zero. A *contraction* (f, g, H) consists of chain maps $f : A \to B$, $g : B \to A$, and a chain homotopy $H : A \to A[1]$ such that

1. $f \circ g = Id_B$,
2. $d_A H + H d_A = g \circ f - Id_A$.

Let A be a C_∞-algebra with maps $m_n^A : A[-1]^{\otimes n} \to A[-1]$. Then B can be given a C_∞-algebra structure such that g can be resolved to a C_∞-algebra map. We describe the transfer map using trees. Let T be a planar tree with n inputs and one output.



FIGURE 3.1. The tree above defines a map $m_n^T : B^{\otimes 7} \to B$ given by $f \circ m_3(Id \otimes H \otimes H) \circ (Id \otimes m_4 \otimes m_2) \circ g^{\otimes 7}$.

Associate to the input edges the map g and the output edge the map f. To each interior vertex of valence k associate the map m_k^A. To each interior edge associate the map H. Then T defines a map $m_n^T : B[-1]^{\otimes n} \to B[-1]$. We refer the reader to Figure 3.1 for an example of a map defined by a planar tree. Then m_n^B is defined as the sum

$$m_n^B = \sum_{\substack{\text{trees } T \\ \text{with n inputs}}} m_n^T.$$

We can define maps $G_n : B[-1]^{\otimes n} \to A[-1]$ in the same way as m_n^B, keeping all the same association of edges and vertices to maps, except associating the output edge with the homotopy H.

Theorem 3.1 ([CG], Theorem 12). *Let (A, m_n^A) be a C_∞-algebra and (f, g, H) be a contraction between A and B. Then (B, m_n^B) is a C_∞-algebra and $\{G_n : B[-1]^{\otimes n} \to A[-1]\}$ is a C_∞-algebra morphism.*

This method of transferring structure works for A_∞ and C_∞-coalgebras as well. The only difference is that trees with one input and n outputs are considered. Suppose $\{c_n : A[-1] \to A[-1]^{\otimes n}\}$ is an A_∞-coalgebra structure on A. Then $c_n^B : B[-1] \to B[-1]^{\otimes n}$ is defined as the sum of all maps c_n^T, where T is a tree with one input and n outputs.

Given a chain complex A with domain $\{A_i\}$ and a dg operad P, a partial P-algebra is then a map of dg operads $P \to \underline{End}(A)$. A partial A_∞-algebra structure on A with domain $\{A_n\}$ is then given by maps $m_n : A_n \to A$ satisfying the same compatibility relations.

We want to transfer partial A_∞-algebra structure by a contraction between A and B using the same methods described in this section. The definition of a domain allows us to do just that. To a corolla with i incoming edges and j outgoing edges, we associate a map $A_i \to A_j$. A general tree is obtained by grafting an outgoing edge of one corolla with an incoming edge of another corolla. We use the ${}_i\circ_j$ compositions, whose existence is assumed in the definition of complex to define the partial endomorphism operad $\underline{End}(A)$, to associate to an (n, m)-tree a map $A_n \to A_m$.

3.2. **Transfer of partial $Frob_\infty^0$-bialgebra by a contraction.** By definition of Koszul Duality, $\mathbb{D}(BiLie^0) \to Frob^0$ is a quasi-isomorphism. Then a vector space V

is a $Frob^0_\infty$-algebra if there is a dioperad map $\mathbb{D}(BiLie^0) \to End(V)$. The complex $\mathbb{D}(BiLie^0)$ is given by

$$\overline{BiLie^0}(m,n)^* \xrightarrow{d''} \bigoplus_{\substack{|edge(T)|=1 \\ \text{reduced } (m,n) \text{ trees}}} \overline{BiLie^0}^*(T) \otimes det(T) \xrightarrow{d''} \bigoplus_{\substack{|edge(T)|=2 \\ \text{reduced } (m,n) \text{ trees}}} \overline{BiLie^0}^*(T) \otimes det(T) \xrightarrow{d''} \cdots .$$

The resulting trees have vertices labelled by elements in $BiLie^0$. We can think of the left most term in this complex as (m,n) trees with no internal vertices. The differential is induced by contracting internal edges.

There are several equivalent definitions for C_∞-algebras, we look for an analogous situation with $Frob^0_\infty$-bialgebras. For our purposes, we look for analogous equivalence of the following two definitions of a C_∞-algebra structure on a vector space V:

1. a C_∞-algebra on V is a map of dg operads $\mathbb{D}(Lie) \to End(V)$
2. a C_∞-algebra on V is given by maps $\{m_n : V[-1]^{\otimes n} \to V[-1]\}$, which vanish on the image of the shuffle product $T^c(V[-1]) \otimes T^c(V[-1]) \to T^c(V[-1])$.

The second definition can be stated as saying a C_∞-algebra is an A_∞-algebra with the extra condition that the structure maps m_n vanish on the image of the shuffle product. It is closely related to the map of operads $Lie \hookrightarrow Ass \to Com$, and the fact that Lie and Com are Koszul.

We use the description of the Frobenius dioperad as $Com^{op} \boxdot Com$ to set up the analogous situation of C_∞-algebras in the dioperad setting. Denote the dioperad $Ass^{op} \boxdot Ass$ by $NC\,Frob^0$ since algebras over $Ass^{op} \boxdot Ass$ are a kind of non-commutative Frobenius bialgebra. A vector space V is an $NC\,Frob^0$ algebra if it has an associative product and a coassociative coproduct, Δ, such that

$$\Delta(ab) = a_{(1)} \otimes a_{(2)}b = ab_{(1)} \otimes b_{(2)}.$$

Denote the dioperad $Ass^{op} \boxdot Ass$ by ϵBi^0, since algebras over this dioperad are infinitesimal bialgebras. A vector space V is an infinitesimal bialgebra, or ϵ bialgebra for short, if it has an associative product and a coassociative coproduct such that

$$\Delta(ab) = a_{(1)} \otimes a_{(2)}b + (-1)^{|a|}ab_{(1)} \otimes b_{(2)}.$$

By Proposition 2.4, $NC\,Frob^0$ and ϵBi^0 are Koszul dual dioperads. A vector space V is an $NC\,Frob^0_\infty$-algebra if there is a dg dioperad map

$$\mathbb{D}(\epsilon Bi^0) \to End(V).$$

An element in $\mathbb{D}(\epsilon Bi^0)$ is represented by a tree whose vertices are labeled by elements in ϵBi^0. But elements in $\epsilon Bi^0 = Ass \boxdot Ass^{op}$ are two level trees whose vertices are labelled by elements in Ass or Ass^{op}. To define a dg dioperad map $\mathbb{D}(\epsilon Bi^0) \to End(V)$, we need to define maps $m^T \in End(V)$ to trees $T \in \mathbb{D}(\epsilon Bi^0)$ which respects the differentials.

In Section 3.1, we showed how an A_∞ or C_∞-algebra can be transferred by a contraction. Then it was shown how to transfer a partial A_∞-algebra. In this section, we discuss how to transfer a $Frob^0_\infty$-bialgebra structure by a contraction.

Let (A_*, d) be a cochain complex, and suppose $\mathbb{D}(\epsilon Bi^0) \to End^0(A_*)$ is a map of dg dioperads. Then given a contraction, with chain maps $f : A_* \to B_*$, $g : B_* \to A_*$, and chain homotopy $H : A_* \to B_*$, we can define a map of dg dioperads $\mathbb{D}(\epsilon Bi^0) \to End^0(B_*)$. Let T whose vertices are labeled by elements in ϵBi^0. Associate the chain map g to the input edges, the chain map f to the output edges, and the chain homotopy H to the interior edges. The map $\mathbb{D}(\epsilon Bi) \to End(A_*)$ associated every element in ϵBi to a map in $End(A_*)$. Associate to each vertex of T labeled by an element in ϵBi the corresponding map in $End(A_*)$. This associates to each tree T Given any $T \in \mathbb{D}(\epsilon Bi^0)$, we get a map $m^T \in End(B_*)$ through this process.

Lemma 3.2. *Let $\mathbb{D}(\epsilon Bi^0) \to End^0(A_*)$ be a map of dg dioperads and (f, g, H) be a contraction between A_* and B_*. Then*

$$\begin{array}{rcl} \mathbb{D}(\epsilon Bi^0) & \to & End^0(B_*) \\ T & \to & m^T \end{array}$$

is a map of dg dioperads.

Proof. We need to show the map respects the differentials. Since the differential for ϵBi^0 is zero, the differential for $\mathbb{D}(\epsilon Bi^0)$ is induced by edge contractions, which we denoted by d''. Then

$$d''(T) = \sum_{T', T''} T' {}_i\circ_j T'',$$

where T' and T'' are trees such that when one edge in $T' {}_i\circ_j T''$ is contracted, we obtain T.

The argument is similar to the one for transferring A_∞-algebras. Consider m^T and take $[m^T, d]$. Using the fact that f and g are chain maps, and $(A, d, \{m_{i,j}\})$ is a $Frob^0_\infty$-bialgebra, we can move the differentials inside the graph, until it meets an interior edge. Then at the interior edge, we have $dH + Hd = g \circ f - Id_A$. The trees with Id_A cancel, because the trees we are summing over compositions of $m^{T'}$ and $m^{T''}$, and A_* is a $Frob^0_\infty$-bialgebra.

We are then left with trees with $g \circ f$. If we split the interior edge into two trees, with the map f associated to the upper tree T' and g associated to the lower tree T'', we see that the upper tree is exactly $m^{T'}$ and the lower tree is $m^{T''}$. So the map respects the differentials. □

Using the arguments in [CG], which showed that when $(A_*, \{m_n\})$ is a C_∞-algebra, the transferred structure on B_* is a C_∞-algebra, we have the corollary.

Corollary 3.3. *If $(A_*, \{m_{i,j}\})$ is a $Frob^0_\infty$-bialgebra, then $(B_*, \{m^B_{i,j}\})$ is a $Frob^0_\infty$-bialgebra.*

The argument in Lemma 3.2 applies to partial structures as well, provided that we can associate a tree to a map on the domain of the partial $Frob^0_\infty$-algebra. This will be the case when $A_* = \Omega^*(M)$ for $A_i = \Omega^*(M^{\times i})$ and $A_* = Curr_*(M)$ and $A_i = Curr_*(M^{\times i})$. We need to make use of kernels and the Dirac delta distributions to define the maps.

4. Topological conformal field theories

Let M be a compact Riemannian manifold. Then $\Omega^*(M)$ is a differential graded commutative algebra, so it is a C_∞-algebra. Let $Har^*(M)$ be the harmonic forms. We show there is a contraction (p, inc, H) where $inc : Har^*(M) \hookrightarrow \Omega^*(M)$ is the inclusion and $p : \Omega^*(M) \to Har^*(M)$ is the projection onto the harmonic forms using the Hodge decomposition.

We have the following operations

$$\begin{aligned} d:\Omega^*(M) &\to \Omega^{*+1}(M)\\ d^*:\Omega^*(M) &\to \Omega^{*-1}(M)\\ \Delta:\Omega^*(M) &\to \Omega^*(M), \end{aligned}$$

where $\Delta = [d, d^*]$. There is also an inner product $\langle\alpha,\beta\rangle = \int \alpha \wedge *\beta$. We can express $e^{-t\Delta}$ as an integral

$$e^{-t\Delta}\alpha(y) = \int K_t(x,y)\alpha(y),$$

where $K_t(x,y) \in \Omega^*(M \times M \times \mathbb{R}^+)$. The operator $e^{-t\Delta}$ is called the heat operator and $K_t(x,y)$ is called the heat kernel.

Definition 4.1. A **heat kernel** for Δ is an element

$$K \in C^\infty(\mathbb{R}^+) \otimes \Omega^*(M) \otimes \Omega^*(M)$$

such that the convolution operator $\widehat{K}_t : \Omega^*(M) \to \Omega^*(M)$ behaves like $e^{-t\Delta}$ where t is the coordinate on $\mathbb{R}^+$. This means that

$$\begin{aligned} \frac{d}{dt}K_t\alpha &= -\Delta K_t\alpha\\ \lim_{t\to 0} K_t\alpha &= \alpha. \end{aligned}$$

When $t = 0$, $e^{-t\Delta}$ is the identity and at $t = \infty$, $e^{-t\Delta}$ is the projection onto the harmonic forms. The chain map $\widehat{K}_t$ is chain homotopic to the identity. Let $L_t = d_x^* K_t$. Then the chain homotopy between $\widehat{K}_t$ is an operator whose kernel is given by

$$\int_0^t L_t.$$

Lemma 4.2. *Let A and B be operators given by kernels K and $K' \in \Omega^*(M \times M)$. Then,*

1. *A is a chain map if and only if K is a closed form,*
2. *A and B are chain homotopic if and only if K and K' are cohomologous.*

Proof. Suppose A is a chain map. Then

$$\begin{aligned} [A,d] &= 0\\ d\int K(x,y)\alpha(y) - \int K(x,y)(d\alpha)(x) &= 0\\ \int (dK(x,y)\alpha(y) + \int K(x,y)(d\alpha)(x) - \int K(x,y)(d\alpha)(x) &= 0\\ \int (dK(x,y))\alpha(y) &= 0, \end{aligned}$$

so $dK(x,y) = 0$. We can reverse this to get the converse.

To prove the second part, it is enough to consider the case when A is null-homotopic. Suppose $K(x,y)$ is exact. Then there is a form J, such that $(d_x + d_y)J = K$. Let J be the kernel of some operator C. We need to show that $[d, C] = A$.

Since X is closed, Stokes's theorem implies $\int_{y\in X} d_y(J(x,y)\alpha)(y) = 0$. This is rewritten as

$$\int (d_y J(x,y))\alpha(y) = \int J(x,y) d_y \alpha(y).$$

Since $(d_y + d_x)J = K$, we see that

$$\begin{aligned} \int d_yJ(x,y)\alpha(y) + \int d_xJ(x,y)\alpha(y) &= \int K(x,y)\alpha(y) \\ \int J(x,y)d_y\alpha(y) + \int d_xJ(x,y)\alpha(y) &= A\alpha \\ C(d\alpha) + dC\alpha &= A\alpha, \end{aligned}$$

which is what we wanted. Again, the converse statement is proved by reversing the order of equalities.

□

The natural setting for this work is on currents. Let $Curr_*(M)$ be the linear dual of $\Omega^*(M)$. We will define a one parameter family of products and coproducts in which $\Omega^*(M)$ forms a sub-object.

Given $\alpha \in \Omega^*(M)$, let $[\alpha]$ be the current $[\alpha]\beta = \int_M \alpha\wedge\beta$. This defines an inclusion $\Omega^*(M) \hookrightarrow Curr_*(M)$. We say $[\alpha]$ is a smooth current.

Kernels can be used to define maps on currents. Let $K^i(x_1, \cdots, x_i)$ be an element in $\Omega^*(M^{\times i})$. Given a current C, let

$$\widehat{K}^{i,j}(C) = [C(x_j)K^i(x_1, \cdots, x_i)].$$

So $\widehat{K}^{i,j}(C)$ is a smooth current in $Curr_*(M^{\times i-1})$. The representing form is obtained by applying C to the j^{th} factor of $K^i \in \Omega^*(M^{\times i})$. Note that if $C = [\alpha]$, then $\widehat{K}^i[\alpha] = [\widehat{K}^i(\alpha)]$, so that the definition of $\widehat{K}^i$ on smooth currents agrees with the definition of $\widehat{K}^i$ on forms.

As they did on $\Omega^*(M)$, K_t and L_t define maps on $Curr_*(M)$. We keep the same notation, $\widehat{K}_t, \widehat{L}_t : Curr_*(M) \to Curr_*(M)$, and $d\widehat{L}_t + \widehat{L}_t d = \widehat{K}_t - Id_{Curr}$.

4.1. **Cellular differential forms.** We now define a form on the moduli space of metrised ribbon graphs. Let $\Gamma(m,n)$ be the set of graphs with m incoming vertices and n outgoing vertices, and let $\gamma \in \Gamma(m,n)$. We will use T to denote a tree in $\Gamma(m,n)$. An edge of γ is external if one of its vertices are external and is internal otherwise. To each edge e of γ, we assign a length $l(e) \in \mathbb{R}^+$. We impose the condition that a path which starts and ends at different outgoing external vertices has positive length. Denote the possible metrics on γ by $Met(\gamma)$. Note that $Met(\gamma)$ is $(\mathbb{R}^+)^{|e(\gamma)|}$ where $|e(\gamma)|$ is the number of edges of γ.

For each internal edge e of γ there is a map

$$Met(\gamma/e) \hookrightarrow Met(\gamma),$$

whose image is the set of all metrics on γ that take the value zero on e.

Definition 4.3. A **cellular differential form** $\omega \in \Omega^i_{cell}(\Gamma(m,n))$ assigns to each graph $\gamma \in \Gamma(m,n)$ a form $\omega_\gamma \in \Omega^i(Met(\gamma), \mathbb{C})$ such that

1. ω_γ is $Aut(\gamma)$ equivariant
2. for each internal edge e,

$$\omega_{\gamma/e} = \omega_\gamma|_{Met(\gamma/e)}.$$

There is an integration pairing

$$\int : C_i(\Gamma(m,n)) \otimes \Omega^i_{cell}(\Gamma(m,n)) \to \mathbb{C}. \tag{4.1}$$

We view $\Gamma(m,n)$ as a topological symmetric monoidal category as follows. The obejcts are the non-negative integers. The morphisms come from the continuous gluing maps

$$\Gamma(l,m)\times\Gamma(m,n)\to\Gamma(l,n).$$

And the monoidal structure comes from disjoint union,

$$\Gamma(m,n)\times\Gamma(s,t)\to\Gamma(m+s,n+t).$$

We wish to consider chains on $\Gamma(m,n)$ with coefficients in a local system, *det*. For $\gamma\in\Gamma(m,n)$, let $det(\gamma)$ be the relative cohomology group of γ,

$$det(\gamma)=\Pi^{n-\chi(\gamma)}det(H^*(\gamma,[n])).$$

With this background, we can define a form. Consider $\alpha=\alpha_1\times\cdots\times\alpha_m\in\Omega^*(M^{\times m})$. For each edge e, let $\omega_e\in\Omega^*(Met(\gamma))\otimes\Omega^*(M\times M)$ be

$$\omega_e=K_{l(e)}+dl(e)L_{l(e)}$$

if e is not an incoming external edge, and

$$(-1)^{p(\Omega^*(M))|\alpha|}(K_{l(e)}+dl(e)L_{l(e)})\alpha_i$$

if e is an incoming external edge with external vertex i. Here, $p(\Omega^*(M))$ is the parity of the map $\Omega^*(M)\to Densities(M)$.

To each edge e there is the two element set of half-edges $H(e)$. Furthermore, since the edges are directed, there is an isomorphism from $H(e)$ to $\{0,1\}$. This allows us to view ω_e as an element in $\Omega^*(Met(\gamma))\otimes\Omega^*(M^{\times H(e)})$. Let $H(\gamma)$ be the set of half edges of γ. We denote the tensor product of all ω_e to get

$$\widetilde{K}_\gamma(\alpha)=\otimes_{e\in E(\gamma)}\omega_e\in\Omega^*(Met(\gamma))\otimes\Omega^*(M^{\times H(\gamma)}).$$

For each vertex $v\in\gamma$, let $H_v(\gamma)$ be the set of half edges at γ. Since there is a cyclic order of the half edges, we can multiply the forms at the half-edges and integrate. This defines a map

$$Tr_v:\Omega^*(M^{\times H_v(\gamma)})\to\Omega^*(M)\xrightarrow{\int_M}\mathbb{C}.$$

Applying Tr_v at each vertex $v\in V(\gamma)\setminus\{\text{outgoing vertex}\}$, then, defines an element

$$K_\gamma(\alpha)=\otimes_{v\in V(\gamma)\setminus\{v_{out}\}}Tr_v\widetilde{K}_\gamma(\alpha).$$

Lemma 4.4 ([Cos], Lemma 4.5.1). *$K_\gamma(\alpha)$ is non-singular. There is a map*

$$K_\gamma:\Omega^*(M)^{\otimes n}\to\Omega^*(Met(\gamma))\otimes\Omega^*(M)^{\otimes m}$$

that commutes with the differentials.

Lemma 4.5 ([Cos], Lemma 4.5.2). *For all $\alpha\in\Omega^*(M^{\times m})$,*

$$K_\gamma(\alpha)|_{Met(\gamma/e)}=K_{\gamma/e}(\alpha).$$

Theorem 4.6 ([Cos], Theorem 4.5.4). *There is a symmetric monoidal functor $C_*(\Gamma,det^{p(\Omega^*(M))})\to Comp_{\mathbb{C}}$ sending*

$$\begin{array}{rcl} n & \mapsto & \Omega^*(M^{\times n}) \\ c & \mapsto & \displaystyle\int_c K(\alpha), \end{array}$$

where c is a chain in $C_*(\Gamma(m,n),det^{p(\Omega^*(M))})$ and $\int_c$ is the integration pairing between $C_*(\Gamma(m,n))$ and $\Omega^*_{cell}(\Gamma(m,n))$.



FIGURE 4.1. The subspace of the moduli space $\Gamma(3,1)$, where the metrics take the values in $[0,t]$ on the internal edge, can be identified with the interval $[-t,t]$. The metrised $(3,1)$ trees associated with the points $-t$, 0, and t in $[-t,t]$ are drawn above.

4.2. **Subspaces of the moduli space.** We consider certain subpsaces of the moduli space of metrised graphs and the cellular differential forms K_γ of the graphs in this space. Associating the graphs with maps on $\Omega^*(M)$ and integrating over the subspaces will define the Frobenius bialgebra we need.

For example, consider the subspace of trees, i.e. genus zero graphs, with n inputs and one output. Furthermore, consider the metrics which take fixed values on the inputs and outputs, but varying between $[0,t]$ on the interior edges. For $n=3$, there are three graphs and the moduli space of metrics is a one parameter family. The form K_T, where $T \in \Gamma(3,1)$ in our subspace, is exact, since it is a one form on $\mathbb{R}$. The moduli space $\Gamma(3,1)$ is described as the interval $[-t,t]$, where $s \in [-t,t]$ represents the metric which takes the value s on the interior edge of the binary tree, and where 0 represents the three corolla. See Figure 4.1 for a description. For $n=4$, the moduli space is a pentagon. Continuing in this way, we obtain cells of the Stasheff associahedron.

We obtain a C_∞-algebra structure on $\Omega^*(M)$ by taking the cellular differential form K_γ and integrating over the above cells in the moduli space. Integrating over these cells define maps m_n^t, for each n, given by

$$\begin{aligned} m_n^t : \Omega^*(M)^{\otimes n} &\to \Omega^*(M) \\ \alpha = \alpha_1 \otimes \cdots \otimes \alpha_n &\mapsto \int K_\gamma(\alpha). \end{aligned}$$

The map $m_2^t(\alpha,\beta) = \widehat{K}_t\alpha \wedge \widehat{K}_t\beta$ is strictly commutative but associative only up to homotopy. The homotopy is given by

$$m_3^t(\alpha,\beta,\gamma) = m_2^t(\widehat{L}_t m_2^t(\alpha,\beta),\gamma) + m_2^t(\alpha,\widehat{L}_t m_2^t(\beta,\gamma)).$$

The definitions of the higher maps proceed in much the same many as transfer of structure described in Section 3.1, where the contraction is given by $f = \widehat{K}_t$, $g = Id$, and chain homotopy $\widehat{L}_t$.

The form K_γ is exact, and so we can use Stokes's theorem to describe the boundary of m_n^t. The boundary consists of integrating over cells of one lower dimension, which can be described as compositions of m_i^t and m_j^t, where $i+j-1=n$. These are precisely the relations for an A_∞-algebra.

5. CONSTRUCTING A PARTIAL $Frob_\infty^0$-BIALGEBRA ON $Curr_*(M)$

5.1. **Double forms $\Omega^*(M \times M)$.** We will briefly review some topics on $\Omega^*(M\times M)$. The reader can find the material in [DR]. A differential form $\gamma \in \Omega^*(M \times M)$ can be viewed as a differential form in $\Omega^*(M)$ with values in $\Omega^*(M)$. That is, given local coordinates $(x_1, \cdots, x_n)$ for $U \subset M$ and $(y_1, \cdots, y_n)$ for $U' \in M$, γ can be expressed as the sum

$$\sum \beta_{i_1 \cdots i_p} dx^{i_1} \wedge \cdots \wedge dx^{i_p},$$

where $\beta_{i_1 \cdots I_p}$ is a form that can be expressed as the sum

$$\sum c_{j_1 \cdots j_q} dy^{j_1} \wedge \cdots \wedge dy^{j_1},$$

where $c_{j_1 \cdots j_q}$ are real numbers. Then γ is a $p+q$ form, and can be expressed by

$$\gamma(x, y) = \sum c_{i_1 \cdots i_p j_1 \cdots j_q} (dx^{i_1} \wedge \cdots \wedge dx^{i_p})(dy^{j_1} \wedge \cdots \wedge dj^{j_q}).$$

The wedge product $\gamma(x, y) \wedge \gamma'(x, y)$ is defined in the obvious way, multiplying the dx terms and dy terms. We can also define the wedge product $\gamma(x, y) \wedge \alpha(x)$, where $\alpha(x) \in \Omega^*(M)$, where we use ony the multiplication in the dx coordinates. Given $\alpha(x)$ and $\beta(y)$, we have the double form $\alpha(x)\beta(y) \in \Omega^*(M \times M)$, which we refer to as the cartesian product of the forms.

In $\Omega^*(M \times M)$, we can take the differential in a single coordinate. We denote it d_x or d_y, and we have the relations

$$d_x^2 = d_y^2 = 0$$
$$d_x d_y = d_y d_x.$$

Similarly, given a chain in one of the coordinates, we can integrate the form in that coordinate, $\int_{x \in c} \alpha(x, y)$, which defines a form in $\Omega^*(M)$.

5.2. **Domain for $Curr_*(M)$.** To define a partial structure on $Curr_*(M)$, we need to specify a domain first (defined in Section 3.1). Consider $\{Curr_*(M^{\times i})\}$. Note that $Curr_*(M)^{\otimes i} \hookrightarrow Curr_*(M^{\times i})$ is a quasi-isomorphism, since over field coefficients, $H_*(M)^{\otimes i} \cong H_*(M^{\times i})$. There is also an $\mathbb{S}_i$ module action, induced by permuting the factors $M^{\times i}$.

We need to define compositions of maps $Curr_*(M^{\times i}) \to Curr_*(M^{\times j})$, for various $i, j > 0$. This is achieved using kernels $\Omega^*(M^{\times n})$ and the Dirac delta distribution, denoted $\widehat{K}_0$. A distribution is a 0-current, i.e. the linear dual of smooth functions. It defines a map on forms by taking a local representation for a form and acting the distribution on the coordinate function. Currents are distribution-valued forms. Since distributions, in general, cannot be multiplied, they do not define maps on currents. We must be careful when using the Dirac delta distribution to define maps and make sure it is only applied to smooth distributions.

Given maps $m_{i_1,1} : Curr_*(M^{\times i_1}) \to Curr_*(M)$ and $m_{i_2,1} : Curr_*(M^{\times i_2}) \to Curr_*(M)$, we define a map $Curr_*(M^{i_1+i_2-1}) \to Curr_*(M)$ as follows. Apply the map m_{i_1} on the factors of $Curr_*(M^{\times i_1+i_2-1})$ as given in the composition and apply $\widehat{K}_t$ to the remaining factors. This will give us an element in $Curr_*(M^{\times i_2})$ to which we apply m_{i_2}. Note that the condition $t > 0$ is necessary, since K_0 is a distribution and cannot be applied to an arbitrary current. Since $\widehat{K}_t$ is chain homotopic to the identity, this change is manageable for our work constructing $Frob_infty^0$-bialgebras.

Similarly, we can define compositions of maps m_{1,j_1}, m_{1,j_2}. In our case, the maps $m_{1,i}$ will land in smooth currents, so we can apply the Dirac delta distribution to the factors.

To define the $\underline{Frob}^0_\infty$-bialgebra, we will have maps $m_{i,j} : Curr_*(M^{\times i}) \to Curr_*(M^{\times j})$. These maps will be compositions of maps of the form $m_{i,1}$, $m_{1,j}$ and $\widehat{L}_t$. The definition of composition extends accordingly.

5.3. **Partial $Frob^0_\infty$-bialgebra structure on $Curr_*(M)$.** To define a $\underline{Frob}^0_\infty$-bialgebra structure on $Curr_*(M)$, we need to define a dg dioperad map

$$\mathbb{D}(BiLie^0) \to \underline{End}^0(Curr_*(M)).$$

We first define the map on $\epsilon Bi^0 \subset \mathbb{D}(\epsilon Bi^0)$. Recall that ϵBi^0 is the infinitesimal bialgebra dioperad and its Koszul dual dioperad is $NC\,Frob^0$. Since $\mathbb{D}(\epsilon Bi^0)$ is a free dioperad, the map on ϵBi^0 extends to $\mathbb{D}(\epsilon Bi^0)$. We then show that the map respects the differentials. Since ϵBi^0 and $NC\,Frob^0$ are Koszul, the map $\mathbb{D}(\epsilon Bi^0) \to \underline{End}^0(Curr_*(M))$ defines a non-commutative $\underline{Frob}^0_\infty$-bialgebra. We then show that the maps are commutative and cocommutative, so that the maps define a $\underline{Frob}^0_\infty$-bialgebra on $Curr_*(M)$.

The process can be viewed as a transfer of structure from $Curr_*(M)$ to itself by the contraction $\widehat{K}_t : Curr_*(M) \to Curr_*(M)$, the identity map $Curr_*(M)$, and homotopy $\widehat{L}_t : Curr_*(M) \to Curr_*(M)$. There is a coproduct given by the diagonal map $\Delta : Curr_*(M) \to Curr_*(M \times M)$. From Section 3, we can transfer the structure to define a C_∞-coalgebra on $Curr_*(M)$. Similarly, we can use the fact that $Im(K_t) \subset \Omega^*(M)$, and use the wedge product to obtain a C_∞-algebra on $Curr_*(M)$. Carrying out this process involves considering binary trees with one input and j outputs for the coalgebra structure and binary trees with i inputs and one output for the algebra structure. Each such tree defines a map in $\underline{End}^0(Curr_*(M))$. These define dg operad maps.

$$\mathbb{D}(Lie) \to \underline{End}^0(Curr_*(M)) \tag{5.1}$$

$$\mathbb{D}(Lie^{op}) \to \underline{End}^0(Curr_*(M)). \tag{5.2}$$

We then use these to define

$$\mathbb{D}(BiLie) \to \underline{End}^0(Curr_*(M)).$$

The process starts by considering binary trees.

Let T be a tree binary tree with i inputs and j outputs. To each such tree, there is a map

$$m^{T,t}_{i,j} : Curr_*(M^{\times i}) \to Curr_*(M^{\times j}),$$

obtained by identifying the input and output edges with $\widehat{K}_t$, the vertices with $\wedge$ or Δ_t as determined by the valence, the internal edges with $\widehat{L}_t$.

Before showing that the maps $m^{T,t}_{i,j}$ defines a $\underline{Frob}^0_\infty$-bialgebra, we consider the cases when i or j is equal to one. When $j = 1$, we have a partial algebra structure. The multiplication $m^t_{2,1} : Curr_*(M \times M) \to Curr_*(M)$ is obtained as follows. Given a double current $A \times B \in Curr_*(M \times M)$, the heat kernel $\widehat{K}_t \times \widehat{K}_t$ is applied to $A \times B$ to obtained a smooth double current $[\alpha \times \beta]$. Then the representing forms are wedged, so that $m^t_{2,1}(A \times B) = [\alpha \wedge \beta]$. Note this product has degree $-d$, where d is the dimension of M. The higher maps $m^t_{i,1}$ are obtained by a process similar to the one found in Section 3.1, and as such defines a partial A_∞-algebra. Since the wedge product is commutative, Theorem 3.1 implies the maps $\{m^t_{i,1}\}$ define a partial C_∞-algebra. Note that $m^t_{i,1}$ has degree $-(i-1)d + (i-2)$, since $i - 1$ is the number



FIGURE 5.1. The cell in $\Gamma(2,2)$ obtained by considering metrics which take values in $[0,t]$ on the internal edge is identified with the interval.

of times the degree $-d$ operation is applied, and $i-2$ is the number of times $\widehat{L}_t$ is applied (internal edges).

We can continue the reasoning for $m^t_{1,j} : Curr_*(M) \to Curr_*(M^{\times j})$. The map $m^t_{1,2} : Curr_*(M) \to Curr_*(M \times M)$ is obtained by applying the $\Delta_* : Curr_*(M) \to Curr_*(M \times M)$ and then flowing the outputs for time t. The map is a degree zero. The higher maps $m^t_{1,j}$ is defined by compositions of $m^t_{1,2}$ and chain homotopy $\widehat{L}_t$. We then obtain a partial C_∞-coalgebra structure on $Curr_*(M)$. Note that $m^t_{1,j}$ has degree $j-2$, since $j-2$ is the number of interior edges, which counts the number of times $\widehat{L}_t$ is applied.

In general, $m^{T,t}_{i,j}$ is obtained by compositions of $m^t_{2,1}$, $m^t_{1,2}$, and $\widehat{L}_t$. This means that $m^{T,t}_{i,j}$ has degree $-(i-2)d + (i+j-3)$.

We claim that this defines $\underline{Frob}^0_\infty$-bialgebra on $Curr_*(M)$. A general element in $\mathbb{D}(\epsilon Bi^0)$ is represented by a tree with vertices labeled by elements in $\epsilon Bi = Ass \boxdot Ass^{op}$. If the tree T is binary and the vertices are labeled by the identity element in $Ass(2)$ or $Ass^{op}(2)$, then let $m^{T,t} \in \underline{End}^0(Curr_*(M))$ be defined as above. If it is a binary tree, and the vertex is labeled by a transposition, apply the transposition to the inputs. If T is not a binary tree, let $m^{T,t} = 0$.

Theorem 5.1. *For $t > 0$, There is a map of dg dioperads*

$$\begin{array}{rcl} \mathbb{D}(BiLie^0) & \to & \underline{End}^0(Curr_*(M)) \\ T & \mapsto & m^{T,t}. \end{array}$$

Proof. To show that the dioperad map respects the differentials, we view the maps $m^{T,t}$ as obtained by integrating over the moduli space. The map $m^t_{2,2}$ is obtained by considering the binary trees with two inputs and two outputs. There are two such trees, after identifying trees by the equivalence relation in the cobar complex. The two trees have one internal edge, and when the metric takes the value zero, the two trees are identified in the moduli space $\Gamma(2,2)$. The cell is identified with $[-t,t]$ as shown in Figure 5.1. Then $b_{2,2} = \int_{[-t,t]} K_\gamma$. The boundary of this map is is given by $(m_{2,1} \otimes 1) \circ (1 \otimes m_{1,2}) - m_{1,2} \circ m_{2,1}$ as desired.

Because the maps $m^{T,t}$ are defined by compositions of $m^t_{2,1}$, $m^t_{1,2}$, and $\widehat{L}_t$ we can apply an inductive argument to show that the higher relations are satisfied.

□

We obtain a $Frob^0_\infty$-bialgebra modeling $(Curr_*(M), \{m^{T,t}\})$ by considering the contraction $\widehat{K}_\infty : Curr_*(M) \to H_*(M)$, $inc : H_*(M) \to Curr_*(M)$, and $\widehat{L}_\infty$, where the inclusion is given by choosing the harmonic form α, such that the smooth current $[\alpha]$ represents the homology class. Then we can transfer the structure over to $H_*(M)$. This process defines a $Frob^0_\infty$-bialgebra algebra on $H_*(M)$, since $H_*(M^{\times i}) \cong H_*(M)^{\otimes i}$.

For different $s, t > 0$, we have different $\underline{Frob}_\infty$-bialgebras on $Cur_*(M)$. But they induce the same $Frob^0_\infty$-bialgebra on $H_*(M)$. To see this, it suffices to show it for $m^t_{i,1}$ and $m^t_{1,j}$, since the rest of the maps are described in terms of these maps.

Lemma 5.2. *Let $s, t > 0$. Suppose $\{m^t_{i,1}\}$ and $\{m^s_{i,1}\}$ are the C_∞-algebra structures on $H_*(M)$ induced by $(Curr_*(M), \{m^t_{i_1}\})$ and $(Curr_*(M), \{m^t_{i_1}\})$ respectively. The maps $m^t_{i,1}$ and $m^s_{i,1}$ are equal for each i. Similarly, the maps C_∞-coalgebra structure on $H_*(M)$ given by $\{m^t_{1,j}\}$ and $\{m^s_{1,j}\}$ are equal.*

Proof. The definition of $m^t_{i,1}$ and $m^s_{i,1}$ uses the inclusion $H_*(M) \hookrightarrow Curr_*(M)$, so we will be looking at smooth currents $[\alpha]$ where α is a harmonic form. The proof uses the following facts:

1. for $\alpha \in Har^*(M)$, $\widehat{K}_t(\alpha) = \alpha$, since $\alpha \in Ker(\Delta)$ and $\widehat{K}_t = e^{-t\Delta}$,
2. for $\alpha \in Har^*(M)$, $\widehat{L}_t(\alpha) = 0$, since
$$\widehat{L}_t(\alpha) = \int_0^t d^*\widehat{K}_u(x,y)(\alpha)du = \int_0^t d^*(\alpha)du = 0,$$
3. for an exact form $d\alpha$, $\widehat{L}_t(d\alpha)$ is exact, since
$$\widehat{L}_t(d\alpha) = \int_0^t d^*_x K_u(x,y) d_y(\alpha)(y)du = d\left(\int_o^t d^*K_t(x,y)\alpha(y)\right).$$

To show that $m^s_{i,1} : H_*(M)^{\otimes i} \to H_*(M)$ and $m^t_{i,1} : H_*(M)^{\otimes i} \to H_*(M)$ are equal, it suffices to show that the maps associated to the i-corollas are zero for $n > 2$.

Recall that $m^t_{i,1} : Curr_*(M^{\times n}) \to Curr_*(M)$ is defined by taking the sum over all binary trees with i inputs and one output. We start with $\alpha_1, \cdots, \alpha_i \in Har^*(M)$ and apply $\wedge_t$ and $\widehat{L}_t$ in the order prescribed by the binary tree. The first fact implies $\alpha_i \wedge_t \alpha_j = \alpha_i \wedge \alpha_j$. The wedge product is not harmonic, but since it is closed, it can be written as the sum of a harmonic form and exact form. Then the second and third facts imply $\widehat{L}_t(\alpha_i \wedge \alpha_j)$ is exact. So when we do perform the operations associated with the binary tree, we end up with an exact form. In cohomology, this map from $H_*(M)^{\otimes n} \to H_*(M)$ is zero, as desired.

The same reasoning applies to the coalgebra maps. □

5.4. **Sub partial $Frob^0_\infty$-bialgebra on $\Omega^*(M)$.** The maps $m^{T,t} \in \underline{End}^0(Curr_*(M))$ send smooth currents to smooth currents. This means that the smooth currents, $\Omega^*(M) \hookrightarrow Curr_*(M)$, are a sub-$\underline{Frob}^0_\infty$-bialgebra of $Curr_*(M)$. We wish to show a stronger statement. That when we dualize $\Omega^*(M)$ and its $\underline{Frob}^0_\infty$-bialgebra, we obtain the $\underline{Frob}^0_\infty$-bialgebra structure on $Curr_*(M)$.

This statement almost follows by construction. Note that $m^t_{2,1} : \Omega^*(M \times M) \to \Omega^*(M)$ is given by the composition

$$\Omega^*(M \times M) \overset{\widehat{K}_t \times \widehat{K}_t}{\to} \Omega^*(M \times M) \overset{Diag^*}{\to} \Omega^*(M).$$

Restricting to the subset $\Omega^*(M) \otimes \Omega^*(M)$, we see that $m^t_{2,1}(\alpha, \beta) = \widehat{K}_t(\alpha) \wedge \widehat{K}_t(\beta)$. When we take the linear dual of $m^t_{2,1}$, we obtain

$$Curr_*(M) \overset{Diag_*}{\to} Curr_*(M \times M) \overset{\widehat{K}_t \times \widehat{K}_t}{\to} Curr_*(M \times M),$$

which is exactly the definition of $m^t_{1,2} : Curr_*(M) \to Curr_*(M \times M)$. The main work was in defining the partial structures appropriately. It did not seem necessary to define a partial algebra structure on $\Omega^*(M)$, since it is possible to define an honest algebra structure. But this would not allow for an easy comparison to $Curr_*(M)$.

If T is a binary tree with a global flow, let T^* be the same tree but with the opposite flow direction. So if T has i inputs and j outputs, T^* has j inputs and i outputs.

Proposition 5.3. *Let* $(Curr_*(M), \{m^{T,t}\})$ *and* $(\Omega^*(M), \{m^{T,t}_\Omega\})$ *be the* $\underline{Frob}^0_\infty$*-bialgebra structures defined above. Then the linear dual of* $m^{T,t}_\Omega$ *is equal to* $m^{T^*,t}$.

Proof. The proof is immediate from the constructions of the structure maps for the $\underline{Frob}^0_\infty$ algebra structures on $\Omega^*(M)$ and $Curr_*(M)$. Note that $m^{T,t}_\Omega$ is the restriction of $m^{T,t}$ to $\Omega^*(M^{\times i})$, viewing an element in $\Omega^*(M^{\times i})$ as a smooth current. But currents are covariant and differential forms are contravariant. So if we have a map $f_{i,j} : M^{\times i} \to M^{\times j}$, on forms we get an induced map $\Omega(M^{\times i}) \to \Omega^*(M^{\times j})$ and thinking of forms as smooth currents, we get the linear dual $Curr_*(M^{\times j}) \to Curr_*(M^{\times i})$. □

6. Homotopy invariance of the $Frob^0_\infty$-bialgebra

We now have two ways of obtaining a $Frob^0_\infty$-bialgebra on $H_*(M)$ and $H^*(M)$, coming from $Curr_*(M)$ and $\Omega^*(M)$, respectively. We can transfer the $\underline{Frob}^0_\infty$-bialgebra structures on $Curr_*(M)$ and $\Omega^*(M)$ given in the previous section to homology and cohomology by a contraction. The contraction is given by using the heat kernel for $t = \infty$ to project onto the harmonic forms. The chain map in the contraction is $\widehat{K}_\infty : \Omega^*(M) \to H^*(M)$ with chain homology $\widehat{L}_\infty$, and similarly for currents. The transfer of structure was described in Section 3.2.

The second way is to consider the cyclic C_∞-algebra structure on $H^*(M)$. The C_∞-algebra is induced by the wedge product on differential forms ($\wedge$ or equivalently $\wedge_0$ in our notation), and the cyclic structure is given by Poincare Duality. Applying the functor

$$\mathcal{D} : \text{cyclic operads} \to \text{dioperads}$$

to the map cyclic $C_\infty \to End(H^*(M))$ defines a $Frob^0_\infty$-bialgebra on $H^*(M)$, as described in Section 2.4. Since Poincare duality and C_∞-algebra on $H^*(M)$ are homotopy invariants of M, showing that the two minimal models are the same proves the main claim of the paper.

We break the proof up into four statements relating the various algebraic structures on currents and forms.

1. $(Curr_*(M), \{m^{T,t}\})$ is a $Frob^0_\infty$-bialgebra. The differential forms $\Omega^*(M)$ is a sub-algebra of $Curr_*(M)$.
2. The linear dual of $(\Omega^*(M), \{m^{T,t}_\Omega\})$ is $(Curr_*(M), \{m^{T,t}\})$.
3. The $\underline{Frob}^0_\infty$-bialgebra $(\Omega^*(M), \{m^{T,t}_\Omega\})$ induces a $Frob^0_\infty$-bialgebra on $H^*(M)$ and the $\underline{Frob}^0_\infty$-bialgebra on $(Curr_*(M), \{m^{T,t}\},)$ induces $Frob^0_\infty$-bialgebra on $H_*(M)$. These models are dual to each other.
4. The Poincare Duality isomorphism from $H^*(M) \to H_*(M)$ identifies the structure maps $\{m^t_{i,j}\}$ in the minimal model for $H^*(M)$ with the structure

maps for $H_*(M)$. That is, the following diagram commutes

$$\begin{tikzcd}
H^*(M)^{\otimes i} \arrow[r, "m^t_{i,j}"] \arrow[d, "PD^{\otimes i}"'] & H^*(M)^{\otimes j} \arrow[d, "PD^{\otimes j}"'] \\
H_*(M)^{\otimes i} \arrow[r, "m^t_{i,j}"] & H_*(M)^{\otimes j}.
\end{tikzcd}$$

Statement (1) is a summary of the work in Sections 5.3 and 5.4, where we constructed $\underline{Frob}^0_\infty$-bialgebras. Statement (2) is proved in Proposition 5.3. Statement (3) is the same as (2) after passing to minimal models. That is, (3) is a corollary of 5.3. Finally, statement (4) follows from the fact that the cup product on cohomology and intersection product on homology are Poincare dual operations.

Theorem 6.1. *The $Frob^0_\infty$-bialgebra on $H^*(M)$ induced by TCFT on $\Omega^*(M)$ is a homotopy invariant of the manifold.*

Proof. We show that the $Frob^0_\infty$-bialgebra on $H^*(M)$ is induced by the cyclic C_∞-algebra on $H^*(M)$. The C_∞-algebra on $H^*(M)$ is obtained by transferring the commutative algebra structure $(\Omega^*(M), \wedge)$ to $H^*(M)$. The cyclic structure is given by capping with the fundamental class $[M]$. Since the C_∞-algebra and $\cap[M]$ are known to be homotopy invariants, showing that the two $Frob^0_\infty$-bialgebras are the same proves the claim.

Lemma 5.2 shows that the maps $m_i : H^*(M)^{\otimes i} \to H^*(M)$ are equal to the maps $m^t_{i,1} : H^*(M)^{\otimes i} \to H^*(M)$ given by the minimal model for $(\Omega^*(M), \wedge_t)$. We write the maps m_i as $m_{i,1}$.

The dual of $m_{i,1} : H^*(M)^{\otimes i} \to H^*(M)$ is a map $c_{1,i} : H_*(M) \to H_*(M)^{\otimes i}$. By statement (3), $\{c_i\}$ defines the coalgebra maps in the $Frob^0_\infty$-algebra on $H_*(M)$ coming from currents. Using Poincare duality isomorphism between $H^*(M)$ and $H_*(M)$ and the maps $\{c_{1,i} : H_*(M) \to H_*(M)^{\otimes i}\}$, we obtain maps $\{m_{1,i} : H^*(M) \to H^*(M)^{\otimes i}\}$. By statement 4, these maps are the coalgebra maps in the $Frob^0_\infty$-bialgebra on $H^*(M)$.

So far, we have shown that $m_{i,1} = m^t_{i,1}$ and $m_{1,j} = m^t_{1,j}$. We need to show that this is true for general map $m^T \in End^0(H^*(M))$. Recall that $m^{T,t}$ is defined by compositions of $\{m^t_{i,1}\}$ and $\{m^t_{1,j'}\}$. Looking at the definition of the functor $\mathcal{D} : \{\text{cyclic operads}\} \to \{\text{dioperads}\}$, we see that the same is true for for $m_{i,j}$. That is, $m_{i,j}$ is also defined by compositions of $\{m_{i',1}\}$ and $\{m_{1,j'}\}$. So we see that the two $Frob^0_\infty$-bialgebras on $H^*(M)$ are the same.

□

References

[A] L. Abrams, Two Dimensional Topological Quantum Field Theories and Frobenius Algebras, *J. Knot Theory Ramifications* (1996), no. 5, 569–587

[DR] G. de Rham, Differentiable Manifolds: Forms Currents, Harmonic Forms, *Springer* (1984)

[CG] X. Cheng, E. Getzler, Transferring homotopy commutative algebraic structures, *J. Pure Appl. Algebra* 212 (2008), no. 11, 2535–2542.

[CL] J Chuang, A. Lazarev, Abstract Hodge Decomposition and Minimal Models for Cyclic Algebras, *Lett Math Phys* (2001), no.89, 33-49

[Cos] K. Costello, Topological Conformal Field Theories and Gauge Theories, *Geometry and Topology* 11 (2007) 1539-1579.

[G] W. L. Gan, Koszul Duality for Dioperads, *Math. Res. Lett.* 10 (2003), no. 1, 109-124.

[1] [CYV] I. Galvez-Carrillo, A. Tonks, B. Vallette, Homotopy Batalin Vilkovisky Algebras, arXiv:0907.2246 , to appear in *J. of Noncommutative Geometry.*

[GK] E. Getzler, M.M Kapranov, Cyclic Operads and Cyclic Homology, *Geometry, topology, & physics* (1995) 167201.

[Gr] J. Granker, Strong homotopy properads, *Int. Math. Res. Not.* 2007, no. 14,

[H] A. Hatcher, Algebraic Topology, Cambridge University Press (2002)

[KS] M. Kontsevich, Y. Soibelman, Homological mirror symmetry and torus fibrations, *Symplectic geometry and mirror symmetry*, (Seoul 2000), World Sci. Publishing, RIver Edge, NJ, 2001, pp. 203-263

[M] M. Markl, Homotopy Algebras are Homotopy Algebras, *Forum Math.* 16 (2004), no. 1, 129160.

[MSS] M. Markl, S. Schnider, J. Stasheff, Operads in Algebra Topology and Physics, Mathematical Surveys and Monographs, vol. 96, *American Mathematical Society*, Providence, RI, (2002)

[MV] S. Merkulov, B. Vallette, Deformation Theory of Representations of Prop(erads) I, *J. Reine Angew. Math.*, 634:51106, 2009

[S] D. Sullivan, Homotopy theory of the master equation package applied to algebra and geometry: a sketch of two interlocking programs, *Banach Center Publ.* 85 (2009), 297-305

[V] B. Vallette, A Koszul Duality for Props, Trans. of Amer. Math. Soc., 359 (2007), 4865-4993

[W] S. Wilson, On the Algebra and Geometry of a Manifold's Chains and Cochains, PhD dissertation, Stony Brook University. August 2005

[W2] S. Wilson, Free frobenius algebra on the differential forms of a manifold arXiv:0710.3550v2

DEPARTMENT OF MATHEMATICS, THE GRADUATE CENTER AT CUNY, 365 FIFTH AVENUE, NEW YORK, NY 10016-4309

E-mail address: `mmiller1@gc.cuny.edu`